European Journal of Anaesthesiology 2002; **19**: iii–ix
© 2002 European Academy of Anaesthesiology
ISSN 0265-0215

Volume Contents

Volume 19 Number 1 January 2002

Editorial

1 Interaction of phenol derivatives with ion channels. *G. Haeseler, M. Leuwer*

Review

9 Anaesthetic agents in paediatric day case surgery: do they affect outcome? *E. W. Moore, B. J. Pollard, R. E. Elliott*

Original articles

18 Sedation assessment in critically ill patients with bispectral index. *M. L. Riess, U. A. Graefe, C. Goeters, H. Van Aken, H. G. Bone*

23 First clinical experience with the rapid-, short-acting amiodarone derivative E 047/1 after cardiac surgery. *H. Gombotz, M. Vicenzi, E. Mahla, P. Rehak, H. Metzler*

32 The effects of fenoldopam on renal function in patients undergoing elective aortic surgery. *M. Halpenny, C. Rushe, P. Breen, A. J. Cunningham, D. Boucher-Hayes, G. D. Shorten*

40 Haemodynamic changes in ischaemic vs. anhepatic pig experimental model of acute liver failure. *K. Theodoraki, G. Kostopanagiotou, V. Smyrniotis, N. Arkadopoulos, A. Prachalias, N. Pyrsopoulos, J. Papadimitriou*

47 The prevention of propofol injection pain by tramadol or ondansetron. *D. Memiş, A. Turan, B. Karamanlioğlu, G. Kaya, Z. Pamukçu*

52 Magnesium infusion reduces perioperative pain. *H. Kara, N. Şahin, V. Ulusan, T. Aydoğdu*

57 Perioperative risk factors in elective pneumonectomy: the impact of excess fluid balance. *A. M. Møller, T. Pedersen, P.-E. Svendsen, A. Engquist*

63 Prophylactic intravenous bolus ephedrine for elective Caesarean section under spinal anaesthesia. *J. P. R. Loughrey, F. Walsh, J. Gardiner*

69 Auditory function following spinal analgesia. Comparison of two spinal needles. *S. K. Malhotra, M. Joshi, S. Grover, S. C. Sharma, A. Datta*

Correspondence

73 Another neurological complication of central venous cannulation. *W. Schummer, C. Schummer*

75 From which countries do chairpersons and invited speakers at important anaesthesia and intensive care meetings come? *A. W. Gelb*

76 Bronchial obstruction treated by fibreoptic-guided suction. *A. A. Klein, C. Thoburn, C. Peters, P. S. G. F. Hardee*

78 Book reviews

81 Guide to Contributors

Volume 19 Number 2 February 2002

Editorial

85 Oasis or mirage? The safety of outpatient dental anaesthesia in hospital. *S. Bricker*

Original articles

88 Inhalation anaesthesia is cost-effective for ambulatory surgery: a clinical comparison with propofol during elective knee arthroscopy. *A. Dolk, R. Cannerfelt, R. E. Anderson, J. Jakobsson*

93 Anaesthetic management and outcome in right-lobe living liver-donor surgery. *G. Cammu, R. Troisi, O. Cuomo, B. de Hemptinne, E. Di Florio, E. Mortier*

99 Effects of clonidine and dopamine on gastric tone. *N.-G. Levein, S.-E. Thörn, M. Watturi*

105 Clinical perception of phantom limb sensation in patients with brachial plexus block. *M. E. Gentili, C. Verton, B. Kinirons, F. Bonnet*

109 Randomized comparison of remifentanil–propofol with a sciatic–femoral nerve block for out-patient knee arthroscopy. *A. Casati, G. Cappelleri, M. Berti, G. Fanelli, P. Di Benedetto, G. Torri*

115 Intraoperative fetal oxygen saturation during Caesarean section: general anaesthesia using sevoflurane with either 100% oxygen or 50% nitrous oxide in oxygen. *R. Parpaglioni, G. Capogna, D. Celleno, P. Fusco*

119 Shorter discharge time after regional or intravenous anaesthesia in combination with laryngeal mask airway compared with balanced anaesthesia with endotracheal intubation. *A. Junger, J. Klasen, B. Hartmann, M. Benson, R. Röhrig, D. Kuhn, G. Hempelmann*

125 Comparison of effects of remifentanil, alfentanil and fentanyl on cardiovascular responses to tracheal intubation in morbidly obese patients. *Z. Salihoğlu, S. Demiroluk, O. Demirkiran, Y. Kose*

129 Postoperative residual curarization with cisatracurium and rocuronium infusions. *G. Cammu, L. de Baerdemaeker, N. den Blauwen, J.-C. de Mey, M. Struys, E. Mortier*
135 *In vitro* effects of 4-chloro-5-ethylphenol in skeletal muscle preparations from malignant hyperthermia susceptible and normal swine. *M. U. Gerbershagen, F. Wappler, M. Fiege, R. Weißborn, P. A. Alberts, F. von Brunnig, J. Schulte am Esch*

Clinical letter
141 Lung image changes during bronchopulmonary lavage estimated using transoesophageal echocardiography. *T. Tsubo, I. Sakai, A. Suzuki, H. Okawa, H. Ishihara, A. Matsuki*

Correspondence
145 Peroperative opening of a patent foramen ovale caused by pulmonary artery obstruction due to knotted pulmonary artery catheter. *R. Tavan, B. Coronel, S. Duperret, P. Mees, R. Blondet*
147 Localized abdominal pain following sympathetic blockade with bretylium for the management of complex regional pain syndrome. *M. Molyneux, R. Venn, J. O'Dwyer*
149 Ultrasound-guided placement of intra-aortic balloon pump. *A. Šustić, I. Medved, O. Šimić*
150 General anaesthesia for emergency surgery during acute organophosphate poisoning. *C. D. A. Goonasekera, C. J. B. Pathiyagoda, C. L. K. Attapattu, M. D. Nagarathne*
152 Postoperative hypoxaemia in a patient with patent foramen ovale. *G. Papadopoulos, S. Sgouropoulou, E. Arnaoutoglou, A. Petrou*
154 Epidural anaesthesia in a patient with POEMS syndrome. *S. Karaca, I. Akgün, A. Agritmis*
155 An unusual case of breathing circuit obstruction. *H. Broculos, P. Sutherland, P. Lanham*

157 Book reviews

160 Author's Checklist

Volume 19 Number 3 March 2002

Editorial
161 End-of-life decisions in The Netherlands. *L. H. D. J. Booij*

Original articles
166 Airway management by physicians wearing anti-chemical warfare gear: comparison between laryngeal mask airway and endotracheal intubation. *Z. Goldik, J. Bornstein, A. Eden, R. Ben-Abraham*
170 Iontophoretically applied lidocaine and the prevention of pain associated with the injection of intravenous propofol: a comparison with intravenous lidocaine. *P. Sharpe, M. Asif, A. Victoria, D. J. Rouchotham*
173 Ketamine and the inhibition of albumin extravasation in chemical peritonitis in rat. *K. Hirota, H. Ishihara, A. Matsuki*
177 Propofol in rats: testing for nonlinear pharmacokinetics and modelling acute tolerance to EEG effects. *H. Ihmsen, A. Tzabazis, M. Schywalsky, H. Schwilden*
189 Subcutaneous epinephrine administration decreases lower oesophageal sphincter pressure and gastro-oesophageal pressure gradient in children under general anaesthesia. *A. Kohjitani, H. Ohara*
193 Alcohol-associated admissions to an adult intensive care unit: an audit. *S. M. Mostafa, B. V. S. Murthy*
197 Bilateral paravertebral somatic nerve block for ventral hernia repair. *Z. Naja, M. F. Ziade, P. A. Lönnqvist*
203 Cuffed oropharyngeal airway as a suitable alternative to the laryngeal mask airway for minor outpatient surgery. *B. Yavaşcaoğlu, H. V. Acar, Ş. F. Kahveci, F. N. Kaya, B. Özcan*

Abstracts
208 Annual Meeting of the European Academy of Anaesthesiology 2001, 28 August–1 September 2001: Graz, Austria.

Correspondence
223 Use of hyperbaric oxygen for hepatic artery thrombosis following adult orthotopic liver transplantation. *T. Dubost, B. Goubaux, M. Dubalde, M. Raucoules-Aime, J. Wolktewicz, J. Gugenheim*
225 Venous gas embolism during liver surgery with argon-enhanced coagulation. *M. L. Ousmane, M. Fleyfel, B. Vallet*
226 Multicompartment block in a parturient undergoing urgent Caesarean section. *M. K. Sanders, M. Y. K. Wee*
229 Minor haemodynamic responses in coronary bypass patients following tracheal intubation using either a lightwand device (Trachlight®) or direct laryngoscopy. *V. Vainionpää, E. Haasvisto, P. Nieminen*

232 Book reviews

Volume 19 Number 4 April 2002

Editorial
237 Ropivacaine and bupivacaine in obstetric analgesia. *G. Capogna*

Review
240 Allergic reactions occurring during anaesthesia. *P. M. Mertes, M.-C. Laxenaire*

Original articles

263 Comparison of bupivacaine 0.2% and ropivacaine 0.2% combined with fentanyl for epidural analgesia during labour. *İ. Aşık, A. Göktuğ, I. Gülay, N. Alkış, A. Uysalel*

271 Fentanyl added to bupivacaine 0.05% or ropivacaine 0.05% in patient-controlled epidural analgesia in labour. *L. Pirbudak, S. Tuncer, H. Koçoğlu, S. Göksu, Ç. Çelik*

276 Effect of VIMA with sevoflurane versus TIVA with propofol or midazolam-sufentanil on the cytokine response during CABG surgery. *S. R. El Azab, P. M. J. Rosseel, J. J. De Lange, E. M. van Wijk, R. van Strik, G. J. Scheffer*

283 Use of sevoflurane during cardiopulmonary bypass decreases incidence of awareness. *B. Celebioğlu, A. G. Pamuk, Ü. Aypar, I. Paşaoğlu*

288 Monitoring intravascular volumes for postoperative volume therapy. *H. Brock, C. Gabriel, D. Bibl, S. Necek*

295 Stability of a sufentanil–ropivacaine mixture in a glass and a PVC reservoir. *G. Brodner, T. Ermert, H. Van Aken, M. Westphal, M. A. E. Marcus, W. Gogarten, C. Goeters, H. Bürkle*

Clinical letter

298 Use of the intubating laryngeal mask airway in a case of ankylosing spondylitis for coronary artery bypass grafting. *J. D'Mello, R. Pagedar, M. Butani, P. Kurkal, K. Pandey*

Correspondence

303 Investigating the usefulness of the laryngeal tube in spontaneously breathing patients. *U. Schultz, W. Bühre*

304 A reply. *D. M. Miller, A. C. Pearce*

305 Efficacy of the laryngeal tube. *T. Asai, I. Hidaka, T. Kubota, S. Kawachi*

306 A reply. *D. M. Miller, A. C. Pearce*

306 Fibreoptic control of the laryngeal tube position. *T. Vollmer, H. V. Genzwuerker, K. Ellinger*

308 Bupivacaine pharmacokinetics and motor blockade following epidural administration of the bupivacaine-sulphobutylether 7-β-cyclodextrin complex in sheep. *J. P. Estebe, C. Ecoffey, G. Dollo, P. Le Corre, F. Chevanne, R. Le Verge*

311 **Book reviews**

Volume 19 Number 5 May 2002

Editorial

313 Cholinesterase estimations revisited: the clinical relevance. *L. Kavalliotis*

Review

317 Central neuraxial blocks and anticoagulation: a review of current trends. *A. Tyagi, A. Bhattacharya*

Original articles

330 Effects of normovolaemic haemodilution on middle cerebral artery blood flow velocity and oxygen delivery. *K. Karadibak, N. Gökmen, S. Erbayraktar, Y. Göktay, A. Taplu, A. Arkan, N. Erkan*

337 Cardiac output assessed by arterial thermodilution during exsanguination and fluid resuscitation: experimental validation against a reference technique. *Z. Friedman, H. Berkenstadt, N. Margalit, E. Segal, A. Perel*

341 Fluoride ion toxicity in rabbit kidney thick ascending limb cells. *M. L. Cittanova, L. Estepa, R. Bourbouze, O. Blanc, M.-C. Verpont, F. Wahbe, P. Cortat, M. Daudon, P. M. Ronco*

350 Comparison of the effects of remifentanil or fentanyl on anaesthetic induction characteristics of propofol, thiopental or etomidate. *W. Wilhelm, A. Budler, A. Huppert, S. Kreuer, O. Bücheler, T. Ziegenfuss, R. Larsen*

357 Delivery performance of commercial target-controlled infusion devices with Diprifusor® module. *S. Schraag, J. Flaschar*

361 Succinylcholine improves the time of performance of a tracheal intubation in prehospital critical care medicine. *A. Ricard-Hibon, C. Chollet, C. Leroy, J. Marty*

368 Intravenous single-dose tramadol versus meperidine for pain relief in renal colic. *O. Eray, Y. Çete, C. Oktay, B. Karoli, S. Akça, N. Çete, F. Eroy*

371 Conditions for insertion of the laryngeal mask airway: comparisons between sevoflurane and propofol using fentanyl as a co-induction agent. A pilot study. *S. B. Gamatra, J. D'Mello, M. Butani, P. Jhamnani*

Clinical letter

376 Aetiology of convulsions due to stellate ganglion block: a review and report of two cases. *A. Mahli, D. Coskun, D. T. Akcali*

Correspondence

381 Monitoring in neuroanaesthesia: update of clinical usefulness. *C. Mélot, J. J. Moraine, J. Berré*

381 Gum elastic bougie, capnography and apnoeic oxygenation. *D. W. Green*

382 Systemic heparinization during postoperative pulmonary embolism induces fatal complications. *N. Kotani, F. Tanioka, T. Tsubo, H. Ishihara, A. Matsuki*

384 Intravenous regional blocks with guanethidine and prilocaine combined with physiotherapy: two children with complex regional pain syndrome, Type 1. *P. P. di Vadi, S. Brill, T. Jack, C. Brown, T. Edwards*

387 **Book reviews**

Volume 19 Number 6 June 2002

Editorial

389 Acid–base management – is it relevant for the study design of hypothermic neuroprotection? *T. Fritsch, A. Piepgras, P. Krafft, S. Schwab, W. Kuschinsky, K. F. Waschke*

Review

395 Pathophysiology of cocaine abuse. *H. A. Brownlow, J. Pappachan*

Original articles

415 Operating room management and strategies in Switzerland: results of a survey. *T. J. Sieber, D. L. Leibundgut*

424 Augmented spontaneous breathing and pulmonary gas exchange during pneumoperitoneum. *A. Loeckinger, C. Hoermann, C. Keller, K. H. Lindner, A. Kleinsasser*

428 Effects of induction of anaesthesia with sufentanil and positive-pressure ventilation on the intra- to extrathoracic volume distribution. *T. von Spiegel, S. Gzanmaris, B. Schoru, M. Scholz, G. J. K. Wietasch, A. Hooft*

436 Psychomotor performance after short-term anaesthesia. *E. Haatvisto, K. Kauranen*

442 Haemodynamic and splanchnic organ blood flow responses during sevoflurane-induced hypotension in dogs. *S. Takeda, N. Sato, T. Tomaru*

447 Predictors of successful neuraxial block: a prospective study. *G. R. de Oliveira Filho, H. P. Gomes, M. H. Z. da Fonseca, J. C. Hoffman, S. G. Pedernieiras, J. H. S. Garcia*

452 Effect of aminophylline on recovery from sevoflurane anaesthesia. *A. Turan, D. Memiş, B. Karamanlioğlu, A. Çolak, Z. Pamukçu, N. Turan*

455 Bacterial contamination of propofol: the effects of temperature and lidocaine. *O. N. Aydin, N. Aydin, B. Gultekin, S. Ozgun, A. Gurel*

Correspondence

459 Differentiating massive carbon dioxide embolism from tension pneumothorax during laparoscopic cholecystectomy. *E. J. Carrero-Cardenal*

460 Ventricular asystole and complete heart block after thoracic epidural analgesia for thymectomy. *T.-C. Liu, C.-H. Hsu, S.-S. Kong, C.-H. Cherng, C.-S. Wong*

463 Book reviews

Volume 19 Number 7 July 2002

Editorial

465 Cardiothoracic anaesthesia and the *European Journal of Anaesthesiology*. *R. Feneck*

Review

466 Laudanosine, an atracurium and cisatracurium metabolite. *V. Fodale, L. B. Santamaria*

Original articles

474 Thiopental impairs neutrophil oxidative response by inhibition of intracellular signalling. *D. Fröhlich, S. Wittmann, G. Rothe, G. Schmitz, K. Taeger*

483 Flurbiprofen does not change the bispectral index and 95% spectral edge frequency during total intravenous anaesthesia with propofol and fentanyl. *K. Hirota, S. Fukushi, S. Baba, A. Matsuki*

487 Acute normovolaemic haemodilution beyond a haematocrit of 25%: ratio of skeletal muscle tissue oxygen tension and cardiac index is not maintained in the healthy dog. *M. Freitag, T. Standl, E.-P. Horn, S. Wilhelm, J. Schulte am Esch*

495 Manipulations in glycogen metabolism and the failure to influence infarct size in the ischaemic rabbit heart. *Y. Gozal, R. A. Wolff, D. M. Van Winkle*

504 Effect of three different doses of urapidil on blood glucose concentrations in the streptozotocin diabetic rat. *K. P. Ittner, M. Bucher, M. Zimmermann, H. F. Grobecker, B. K. Krämer, K. Taeger*

510 Can computers help in assessing children's postoperative pain? Initial validation of a computer-assisted interview. *S. Watson, R. Calam, P. Jimmieson*

517 Comparison of different priming techniques on the onset time and intubating conditions of rocuronium. *B. Yavaşcaoğlu, V. Cebelli, N. Kelebek, N. Uçkunkaya, O. Kutlay*

522 Effects of adding alfentanil or atracurium to lidocaine solution for intravenous regional anaesthesia. *N. Kurt, I. Kurt, B. Aygünes, H. Oral, M. Tulunay*

Correspondence

526 Acute abdomen after coronary artery bypass surgery masked by thoracic epidural analgesia. *Z. Jankovic, D. Radojevic, V. Neskovic, D. Stamenkovic*

528 Epidural anaesthesia for arthroscopic knee surgery in a patient with Takayasu's arteritis. *S. Karaca, I. Akgün*

530 Workforce and regional distribution of anaesthesiologists in Japan. *K. Taga, H. Fujihara, H. Baba, T. Yamakura, K. Shimoji*

532 Insidious hypothyroidism unmasked after operation. *G. Vretzakis, E. Ferdi, B. Papaziogas*

535 Book reviews

537 Guide to Contributors

Volume 19 Number 8 August 2002

Editorial

543 Transcranial cerebral oximetry, transcranial Doppler sonography, and heart rate variability: useful neuromonitoring tools in anaesthesia and intensive care? *G. Schwarz, G. Litscher*

Review

550 Isoprostanes – markers of ischaemia reperfusion injury. *H. Sakamoto, T. B. Corcoran, J. G. Laffey, G. D. Shorten*

Original articles

560 Impact of the reduction of anaesthesia turnover time on operating room efficiency. *E. Sokolovic, P. Biro, P. Wyss, C. Werthemann, U. Haller, D. Spahn, T. Szucs*

564 Lidocaine-induced cell death in a human model of neuronal apoptosis. *P. Friederich, T. P. Schmitz*

571 Voltage-dependent block of neuronal and skeletal muscle sodium channels by thymol and menthol. *G. Haeseler, D. Maue, J. Grosskreutz, J. Bufler, B. Nentwig, S. Piepenbrock, R. Dengler, M. Leuwer*

580 Target-controlled propofol requirements at induction of anaesthesia: effect of remifentanil and midazolam. *D. H. Conway, S. K. Hasan, M. E. Simpson*

585 Using the Sustained Attention to Response Test to detect cognitive dysfunction after day case surgery. *J. P. Thompson, S. C. Rushman, A. J. Fox, A. J. Lloyd, R. A. Atcheson*

589 Anaesthetic and postoperative analgesic effects of spinal clonidine as an additive to prilocaine in the transurethral resection of urinary bladder tumours. *X. Santiveri, A. Arxer, I. Plaja, M. T. Metje, B. Martinez, A. Villalonga, M. López*

594 Spinal block or total intravenous anaesthesia with propofol and remifentanil for gynaecological outpatient procedures. *G. Dawelli, M. Berti, A. Casati, A. Albertin, F. Deni, F. Nobili, G. Torri*

600 Decrease in minimum alveolar concentration of sevoflurane during anaesthesia and arthroscopy. *H. Higuchi, Y. Adachi*

604 Effects of chronic occupational exposure to anaesthetic gases on the rate of neutrophil apoptosis among anaesthetists. *R. Tyther, M. Halligan, J. Wang, H. P. Redmond, G. Shorten*

609 Effects of sodium nitroprusside-induced controlled hypotension on pancreatic function assessed by pancreatitis-associated protein in patients undergoing radical prostatectomy. *S. N. Piper, S. W. Suttner, W. H. Malack, B. Kunle, G. Haisch, J. Boldt*

Correspondence

614 Morbidity in electroconvulsive therapy. *M. L. Meignan*

615 A reply. *E. Tecoult, N. Nathan*

616 Morbidity in electroconvulsive therapy. *B. Wild, G. W. Eschweiler, M. Bartels*

616 Accidental injection of intravenous bupivacaine. *S. Karaca, E. Ö. Ünlüsoy*

618 Book reviews

Volume 19 Number 9 September 2002

Editorial

621 Murder and the ICU. *G. R. Park, S. N. Khan*

Review

624 Cardiac arrest, chain of survival and Utstein style. *A. Gullo*

Original articles

634 Ketamine effects on human neuronal Na^+ channels. *G. Rackstegel, P. Friederich, B. W. Urban*

641 Accumulation of *S*(+) enantiomer in human beings after general anaesthesia with isoflurane racemate. *H. A. Haeberle, H. G. Wahl, H. Jakubetz, H. Krause, R. Schmidt, V. Schurig, H.-J. Dieterich*

647 Comparison between nasal and intravenous desmopressin for the treatment of aminosalicylic acid-induced platelet dysfunction. *S. Schulz-Stübner, D. Zielske, R. Rossaint*

652 Gastric emptying in post-thoracotomy patients receiving a thoracic fentanyl-bupivacaine epidural infusion. *A. Guha, N. D. A. Scaun, S. A. Rogers, S. H. Pennefather, G. N. Russell*

658 Postoperative pain management in orthopaedic patients: no differences in pain score, but improved stress control by epidural anaesthesia. *H. A. Adams, P. Saatweber, C. S. Schmitz, H. Hecker*

666 Epidural combination of ropivacaine with sufentanil for postoperative analgesia after total knee replacement: a pilot study. *S. Kampe, C. Diefenbach, B. Kanis, M. Auweiler, P. Kienicke, K. Cranfield*

672 Epidural morphine injection after combined spinal and epidural anaesthesia. *C. Takenaka-Hamaya, Y. Hamaya, S. Dohi*

677 Propofol in low doses causes redistribution of body heat in male volunteers. *I. Noguchi, T. Matsukawa, M. Ozaki, Y. Amemiya*
682 Posterior subgluteal approach to block the sciatic nerve: description of the technique and initial clinical experiences. *P. Di Benedetto, A. Casati, L. Bertini, G. Fanelli*

Correspondence
687 Anaesthesia for the professional singer. *C. L. Errando*
688 Randomizing patients to permit the development of perioperative hypothermia is inappropriate. *M. G. Booth, G. Gallagher, J. Kinsella*
688 A reply. *J. Lundberg, K. Persson*
689 Dilated cardiomyopathy in dystrophic epidermolysis bullosa: a lethal complication of epidermolysis bullosa. *P. M. D. Cunnington, R. Addison*
690 Simple preoperative assessment to reduce the risk of traumatic epistaxis during nasotracheal intubation. *R. Kayarkar, T. J. Woolford, G. A. Francis*
691 Anaphylactic reaction after cisatracurium administration. *E. Iannuzzi, M. Iannuzzi, M. S. Pedicini, V. Civillo, M. Chiefari, G. Sacerdoti*
693 Transient respiratory compromise after infraclavicular vertical brachial plexus blockade. *F. M. Heid, T. Kern, A. M. Brambrink*

695 Book reviews

Volume 19 Number 10 October 2002

Editorial
701 The Cochrane 1998 Albumin Review – not all it was cracked up to be. *P. J. Horsey*

Review
705 Neurological complications following central neuraxial blocks: are there predictive factors? *O. De Tommaso, A. Caporuscio, V. Tagariello*

Original articles
717 Auditory- and somatosensory-evoked potentials in cerebral malaria and anaesthesia: a comparison. *C. Thornton, R. S. Heyderman, M. Thornley, N. Curtis, J. Madke, G. Pasvol, D. E. F. Newton*
727 Cost-reduction analysis of propofol versus sevoflurane: manitenance of anaesthesia for gynaecological surgery using the bispectral index. *M. M. R. F. Struys, A. A. L. Somers, N. Van Den Eynde, L. L. G. Herregods, D. Dyzers, H. M. Robays, E. P. Mortier*
735 Application of semilinear canonical correlation to the measurement of the electroencephalographic effects of volatile anaesthetics. *J. Bruhn, B. Rehberg, H. Röpcke, T. Bouillon, A. Hoeft*
742 Anaesthesia for advanced rectal cancer patients treated with combined major resections and intraoperative radiotherapy. *G. H. H. Mavmaerts, A. A. J. Van Zundert, V. C. H. Meeuwen, H. Martijn, H. J. T. Rutten*
749 QT interval and QT dispersion during the induction of anaesthesia in patients with subarachnoid haemorrhage: a comparison of thiopental and propofol. *P. E. Tanskanen, J. V. Kyttä, T. T. Randell*
755 Anaesthesia manpower in Belgium until 2020: nurse anaesthetists – a possible solution? *J. L. Demeere*

Correspondence
760 Sigmoid capnometry in abdominal aortic aneurysm surgery. *M. Björck*
761 A reply. *G. Lebuffe, B. Vallet*
762 Oasis or mirage? The safety of outpatient dental anaesthesia in hospital. *J. A. W. Wildsmith*
763 A reply. *S. Bricker*
764 Difficulty in use and device failure with the intubating laryngeal mask airway®. *G. Caponas, M. Popat*
766 Salvage with a haemofiltration technique unsuitable in hip prosthesis surgery due to free haemoglobin. *M. Tylman, J.-P. Arnestad, A. Bengtsson, J.-P. Bengtson*
768 Anaesthesia for Caesarean section in the presence of aortic coarctation. *T. Togal, M. Durmus, A. Koroglu, S. Demirbilek, K. Karaaslan, O. Ersoy*
770 Hysteroscopic syndrome. *N. Kayacan, B. Karsli, G. Zorlu, L. Düşemeci, M. Erman*
772 Thoracic epidurals and avoidance of potential neurological damage. *V. Tucker, F. King, J. Nightingale*

774 Book reviews

Volume 19 Number 11 November 2002

Editorial
777 Liver surgery has become simpler. *D. Franco*

Review
780 Anaesthesia for elective liver resection: some points should be revisited. *C. Lentschener, Y. Ozier*

Original articles
789 Doppler indices of diastolic transmitral flow velocity are invalid indicators of myocardial ischaemia during high-dose dobutamine infusion in anaesthetized patients. *M. Filipovic, M. D. Seeberger, R. Rohlfs, O. Dergelm, W. Studer, D. Atar, P. Buser, K. Skarvan*

796 Myocardial ischaemia in patients with impaired left ventricular function undergoing coronary artery bypass grafting – milrinone versus nifedipine. *T. Möllhoff, C. Schmidt, H. Van Aken, E. Berendes, H. Buerkle, P. Marmann, T. Reinbold, R. Prenger-Berninghoff, T. D. T. Tjan, H. H. Scheld, M. C. Deng*

803 Randomized controlled trial investigating the effect of transcervical papaverine and bupivacaine on postoperative analgesia following laparoscopic sterilization. *A. Ng, A. Habib, A. Swami, G. Smith, D. Nunns, A. C. Davidson*

808 Effects of volume and concentration of lidocaine on epidural anaesthesia in pregnant females. *M. Nakayama, J. Yamamoto, H. Ichimoe, S. Yamamoto, N. Kanaya, A. Namiki*

812 Epidural bupivacaine with sufentanil or fentanyl during labour: a randomized, double-blind study. *O. K. Rolfsing, E. Skogvoll, P. C. Borchgrevink*

819 Sufentanil or clonidine for blunting the increase in intraocular pressure during rapid-sequence induction. *M. Georgiou, A. Papalapani, H. Argiriadou, P. Papagiannopoulou, G. Katsikis, E. Kaprini*

823 Effect of different doses of cisatracurium on intraocular pressure in sedated patients. *S. M. Sator-Katzenschlager, M. J. Oehmke, M. Kontaratos, A. Wedrich, G. Heinze, C. Weinstabl*

829 Does ibuprofen increase perioperative blood loss during hip arthroplasty? *R. Slappendel, E. W. G. Weber, B. Benraad, R. Dirksen, M. L. T. Bugter*

Correspondence

832 Central venous catheter malposition in the azygos arch. *W. Schummer, C. Schummer*

834 Sciatic nerve block and the improvement of femoral nerve block analgesia after total knee replacement. *A. Weber, R. Fournier, E. Van Gessel, Z. Gamulin*

836 Anaesthetic approach in a case of Goldenhar's syndrome. *C. Kaymak, Y. Gülhan, A. O. Ozcan, B. Baltaci, N. Unal, M. A. Safak, H. Oguz*

839 Remifentanil and acute intermittent porphyria. *M. Durmus, A. Turkuz, T. Togal, A. Koroglu, H. I. Toprak, M. O. Ersoy*

841 Midodrine, an alternative to intravenous vasopressor therapy after spinal surgery. *B. O'Donnell, A. Synnott*

842 General anaesthesia with remifentanil and propofol for a patient with centronuclear (myotubular) myopathy. *A. Tokarz, T. Gaszyński, W. Gaszyński, P. Arkuszewski*

844 Tongue piercing in an eclamptic patient. *R. W. A. Noucicki, P. T. Bull*

845 Anaesthetic implications of Aicardi's syndrome. *B. O'Brien, D. Louw, M. Awan*

846 Pulmonary oedema after venous air embolism during craniotomy. *C. El Kettani, R. Badaoui, M. Fikri, P. Jeanjean, D. Montpellier, J. Tchamitchian*

848 Spontaneous restoration of decreased systemic vascular resistance after spinal anaesthesia. *M. Kamenik*

851 Book review

Volume 19 Number 12 December 2002

Review

853 Long QT syndrome and anaesthesia. *N. A. Wisely, E. A. Shipton*

Original articles

860 Oxygen transport and myocardial function after the administration of albumin 5%, hydroxyethylstarch 6% and succinylated gelatine 4% to rabbits. *D. G. Himpe, S. G. De Hert, K. M. Vermeyen, H. F. Adriaensen*

868 Preload and haemodynamic assessment during liver transplantation: a comparison between the pulmonary artery catheter and transpulmonary indicator dilution techniques. *G. Della Rocca, M. G. Costa, C. Coccia, L. Pompei, P. Pietropaoli*

876 Haemodynamic and splanchnic organ blood-flow responses to hypotension induced by the calcitonin gene-related peptide and sodium nitroprusside in dogs. *S. Takeda, T. Tomaru*

883 Neuromuscular effects of rapacuronium on the diaphragm and skeletal muscles in anaesthetized patients using cervical magnetic stimulation for stimulating the phrenic nerves. *O. Moerer, C. Baller, J. Hinz, H. Buscher, T. A. Crozier*

888 Fate of abstracts from the Paris 1995 European Society of Anaesthesiologists meeting. *J. Castillo, R. Garcia-Guasch, I. Cifuentes*

Correspondence

894 Is traditional fasting policy changing? Results of a Swiss survey on current practice. *C. K. Hofer, A. Zollinger, M. Tucci, E. van Gessel, E. Ahn, R. Klaghofer*

896 General anaesthesia for a child with Rubinstein-Taybi syndrome. *A. Tokarz, T. Gaszyński, W. Gaszyński, P. Arkuszewski*

897 Anaesthesia for open-heart surgery in a patient with Weaver's syndrome. *B. Celebioğlu, F. Yener*

899 Effects of thiamylal, sevoflurane and isoflurane on the cortically recorded somatosensory evoked potentials. *K. Fukui, T. Moriwaka, K. Hisada, S. Nishio, K. Irita, S. Takahashi*

902 Book review

904 Subject index

908 Author index

Volume contents

Referees for 2002

© 2002 European Academy of Anaesthesiology, *European Journal of Anaesthesiology* 19: iii–ix